

On the number of eigenvalues of the discrete one-dimensional Schrödinger operator with a complex potential

Artem Hulko¹ 

Received: 17 October 2016 / Accepted: 9 November 2016 / Published online: 21 November 2016
© The Author(s) 2016. This article is published with open access at Springerlink.com

Abstract We study the eigenvalues of the discrete Schrödinger operator with a complex potential. We obtain bounds on the total number of eigenvalues in the case where V decays exponentially at infinity.

Keywords Discrete Schrödinger · Complex potential · Total number of eigenvalues

1 Introduction and main results

Let $\mathfrak{H} = \ell^2(\mathbb{Z}_+)$ be the Hilbert space of square summable sequences on $\mathbb{Z}_+ = \{1, 2, 3, \dots\}$. Let $V: \mathfrak{H} \mapsto \mathfrak{H}$ be the operator of multiplication by a bounded complex-valued function on \mathbb{Z}_+ . We study the spectral properties of the Schrödinger operator H , defined in \mathfrak{H} by

$$(Hu)_j = \sum_{|l-j|=1} u_l + V_j u_j, \quad \forall j \geq 2. \quad (1.1)$$

Additionally, we set

$$(Hu)_1 = u_2 + V_1 u_1.$$

Communicated by Ari Laptev.

✉ Artem Hulko
ahulko@uncc.edu

¹ Department of Mathematics and Statistics, University of North Carolina at Charlotte, Charlotte, NC 28223, USA

Note that H is a bounded operator. The spectrum of the self-adjoint operator $H_0 = H - V$ coincides with the interval $[-2, 2]$ and is absolutely continuous. Let λ_j denote the eigenvalues of the operator (1.1). We are interested in an estimate of the total number N of eigenvalues λ_j in the case where the sequence V_j decays exponentially fast.

More precisely, we shall prove the following two theorems:

Theorem 1.1 *The number N of eigenvalues of H in $\ell^2(\mathbb{Z}_+)$, counting algebraic multiplicities, satisfies*

$$N \leq \frac{1}{2 \ln \Lambda} \left(\frac{2\Lambda^2}{\Lambda^2 - 1} \sum_{n=1}^{\infty} \Lambda^{2n} |V_n| \right)^2,$$

for any $\Lambda > 1$.

A similar result for a continuous operator was proved in [13] by Frank, Laptev and Safronov.

We also establish a slightly different estimate:

Theorem 1.2 *The number N of eigenvalues of H in $\ell^2(\mathbb{Z}_+)$, counting algebraic multiplicities, satisfies*

$$N \leq \frac{1}{\ln \Lambda} \frac{\Lambda^2}{(\Lambda^2 - 1)} \left(\sum_{n=1}^{\infty} \Lambda^n |V_n|^{1/2} \right)^2,$$

for any $\Lambda > 1$.

Note that the right hand sides of both estimates can be finite only in the case where V is an exponentially decaying potential. It turns out that N might be finite even in the case when the potential decays slower. For instance, the operator $-d^2/dx^2 + V(x)$ on the half-line $[0, \infty)$ has finitely many eigenvalues if $|V| \leq C \exp(-c\sqrt{x})$ for some $C, c > 0$. This remarkable result was proved by Pavlov in [23]. It was established that the eigenvalues can not accumulate to a point of the positive half-line, which is enough to conclude that the set of all eigenvalues is finite.

On the other hand, there is another remarkable result of Pavlov (see [24]), which says that, for any $0 < p < 1/2$, there exists a complex-valued potential V satisfying $|V| \leq C \exp(-c|x|^p)$ and a complex number θ , such that the operator $-d^2/dx^2 + V(x)$ with the boundary condition $\psi'(0) = \theta\psi(0)$ has infinitely many eigenvalues. Another interesting result was recently established by Bögli [2]. It was shown that there exists a potential for which the eigenvalues accumulate to every point on $[0, \infty)$.

2 Zeroes of analytic functions

The following proposition gives a useful bound on the zeroes of an analytic function in the complement of the disc of radius $R > 0$.

Proposition 2.1 *Let $0 < R < 1$. Let $a(\cdot)$ be an analytic function in $\{k: |k| > R\}$. Assume that $a(\cdot)$ is continuous up to the boundary and satisfies*

$$a(k) = 1 + O(|k|^{-1}) \text{ as } |k| \rightarrow \infty \text{ in } \{k: |k| > R\}. \quad (2.1)$$

Assume also that for some $A \geq 1$,

$$|a(k)| \leq A, \text{ if } |k| = R. \quad (2.2)$$

Then the zeroes k_j of $a(\cdot)$ in $\{k: |k| > R\}$, repeated according to their multiplicities, satisfy

$$\prod_j \left(\frac{|k_j|}{R} \right) \leq A. \quad (2.3)$$

Proof We introduce the Blaschke product

$$B(k) = \prod_j \frac{k - k_j}{R - R^{-1} \bar{k}_j k}.$$

Clearly, $a(k)/B(k)$ is an analytic and non-zero in $\{k: |k| > R\}$. Consequently, $\log(a(k)/B(k))$ exists and is analytic in $\{k: |k| > R\}$. Let C_R denote the circle $\{k \in \mathbb{C}: |k| = R\}$, traversed counterclockwise.

Then, according to the residue calculus,

$$\int_{C_R} \log \frac{a(k)}{B(k)} \frac{dk}{k} = 2\pi i \lim_{k \rightarrow \infty} \log \frac{a(k)}{B(k)} = 2\pi i \sum_j \log \frac{\bar{k}_j}{-R},$$

and therefore

$$\int_{-\pi}^{\pi} \log \frac{a(Re^{i\varphi})}{B(Re^{i\varphi})} d\varphi = 2\pi \sum_j \log \frac{\bar{k}_j}{-R}. \quad (2.4)$$

We note that $|B(Re^{i\varphi})| = 1$ if $\varphi \in \mathbb{R}$ and, therefore,

$$\operatorname{Re} \int_{-\pi}^{\pi} \log \frac{a(Re^{i\varphi})}{B(Re^{i\varphi})} d\varphi = \int_{-\pi}^{\pi} \ln \left| \frac{a(Re^{i\varphi})}{B(Re^{i\varphi})} \right| d\varphi = \int_{-\pi}^{\pi} \ln |a(Re^{i\varphi})| d\varphi. \quad (2.5)$$

On the other hand,

$$\operatorname{Re} \sum_j \log \frac{\bar{k}_j}{-R} = \sum_j \ln \frac{|k_j|}{R}. \quad (2.6)$$

We conclude from (2.4), (2.5) and (2.6) that

$$\int_{-\pi}^{\pi} \ln |a(Re^{i\varphi})| d\varphi = 2\pi \sum_j \ln \frac{|k_j|}{R}. \quad (2.7)$$

Finally, by (2.2),

$$\int_{-\pi}^{\pi} \ln |a(Re^{i\varphi})| d\varphi \leq 2\pi \ln A, \quad (2.8)$$

Inequality (2.3) now follows from (2.7) and (2.8). \square

Corollary 2.2 *Let $0 < R < 1$. Let $a(\cdot)$ be an analytic function in $\{k: |k| > R\}$ satisfying (2.1). Assume that, for any $R' > R$ sufficiently close to R , condition (2.2) holds with R replaced by R' . Then the number*

$$\mathcal{N} := \#\{j: |k_j| \geq 1\}$$

of zeroes k_j of $a(\cdot)$ in $\{k: |k| \geq 1\}$, repeated according to their multiplicities, satisfies

$$\mathcal{N} \leq \frac{\ln A}{\ln 1/R}.$$

Proof We apply Proposition 2.1 for every $R' > R$ sufficiently close to R and obtain

$$\sum_j (\ln |k_j| - \ln R')_+ \leq \ln A$$

Clearly, we have

$$\sum_j (\ln |k_j| - \ln R')_+ \geq |\ln R'| \cdot \#\{j: |k_j| \geq 1\}.$$

Consequently,

$$|\ln R'| \cdot \mathcal{N} \leq \ln A.$$

The corollary follows by passing to the limit $R' \rightarrow R$. \square

3 Classes of compact operators and determinants

Let $1 \leq p < \infty$. We say that a compact operator T belongs to the Schatten class \mathfrak{S}_p if its singular values $s_j(T)$ satisfy

$$\|T\|_{\mathfrak{S}_p}^p := \sum_j s_j^p(T) < \infty.$$

The functional $\|\cdot\|_{\mathfrak{S}_p}$ is the norm on \mathfrak{S}_p .

Let $K \in \mathfrak{S}_n$ with $n \in \mathbb{N}$. Let $\lambda_j(K)$ denote the eigenvalues of K , repeated according to algebraic multiplicities. The n -th order regularized determinant $\det_n(1 + K)$ is defined by

$$\det_n(1 + K) := \prod_j \left((1 + \lambda_j(K)) \exp \left(\sum_{m=1}^{n-1} \frac{(-1)^m}{m} \lambda_j(K)^m \right) \right).$$

The following property is well-known, but we include a proof for the sake of completeness.

Lemma 3.1 *Let $n \in \mathbb{N}$ and let $K \in \mathfrak{S}_n$. Then*

$$\ln |\det_n(1 + K)| \leq \Gamma_n \|K\|_{\mathfrak{S}_n}^n,$$

where Γ_n is a positive constant independent of K . In particular,

$$\Gamma_1 = 1 \quad \text{and} \quad \Gamma_2 = 1/2. \quad (3.1)$$

Proof To prove the lemma, let $f(z) := (1 + z) \exp \left(\sum_{m=1}^{n-1} \frac{(-1)^m}{m} z^m \right)$. Then $\ln |f(z)|$ can be bounded by a constant times $|z|^n$ for small $|z|$ and by a constant times $|z|^{n-1}$ for large $|z|$. Thus, $\ln |f(z)| \leq \Gamma_n |z|^n$, and so

$$\ln |\det_n(1 + K)| \leq \Gamma_n \sum_j |\lambda_j(K)|^n$$

By Weyl's inequality [26, Thm. 1.15], the sum on the right side does not exceed $\|K\|_{\mathfrak{S}_n}^n$. A simple computation shows that for $n = 1$ and $n = 2$ one can take $\Gamma_1 = 1$ and $\Gamma_2 = 1/2$, respectively (see [27]). \square

We now recall the Birman–Schwinger principle. We state it in the case, where H_0 is a bounded self-adjoint operator and $V = G^*G_0$. We will assume that G_0 and G are compact operators. Now, set

$$H = H_0 + V.$$

The Birman–Schwinger principle states that $z \in \rho(H_0)$ is an eigenvalue of H if and only if -1 is an eigenvalue of the Birman–Schwinger operator $G_0(H_0 - z)^{-1}G^*$. Moreover, the corresponding geometric multiplicities coincide.

The following lemma says that even the algebraic multiplicities of eigenvalues of H can be characterizes in terms of a quantity related to the Birman–Schwinger operator.

Lemma 3.2 *Let $n \in \mathbb{N}$. Assume that $G_0(H_0 - \zeta)^{-1}G^* \in \mathfrak{S}_n$ for all $\zeta \in \rho(H_0)$. Then the function $\zeta \mapsto \det_n(1 + G_0(H_0 - \zeta)^{-1}G^*)$ is analytic in $\rho(H_0)$. A point $z \in \rho(H_0)$ is an eigenvalue of H if and only if $\det_n(1 + G_0(H_0 - z)^{-1}G^*) = 0$. Moreover, the order of the zero coincides with the algebraic multiplicity of the corresponding eigenvalue.*

Analyticity of the function $\zeta \mapsto \det_n(1 + G_0(H_0 - \zeta)^{-1}G^*)$ is well-known (see, e.g., [27]), as well as the result about the algebraic multiplicity in the case $n = 1$. The result for the general n is essentially due to [18]; you may also refer to [11] for an extension of the proof to the present setting.

4 Resolvent bounds

In this section we collect trace ideal bounds for the Birman–Schwinger operator

$$K(k) = \sqrt{V}(H_0 - z)^{-1}\sqrt{|V|}, \quad z = k + k^{-1}, \quad |k| \geq 1. \quad (4.1)$$

We use the notation $\sqrt{V(x)} = V(x)/\sqrt{|V(x)|}$ if $V(x) \neq 0$ and $\sqrt{V(x)} = 0$ if $V(x) = 0$.

We remind the reader that $\mathfrak{H} = \ell^2(\mathbb{Z}_+)$, and H_0 in (4.1) denotes the free Jacobi operator on \mathbb{Z}_+ . From the explicit expression of its matrix it is easy to see that, if V has a compact support, then $K(k)$ admits an analytic continuation to $\mathbb{C} \setminus \{0\}$. The following proposition gives a bound on the Hilbert–Schmidt norm.

Proposition 4.1 *For any $k \in \mathbb{C} \setminus \{0\}$ with $|k| < 1$,*

$$\|K(k)\|_{\mathfrak{S}_2} \leq \frac{2}{1 - |k|^2} \sum_{n=1}^{\infty} |k|^{-2n} |V_n|,$$

Proof The matrix of $(H_0 - z)^{-1}$ is given by

$$g_k(n, m) = \frac{k}{k^2 - 1} \left(k^{-|n-m|} - k^{-(n+m)} \right),$$

which satisfies

$$|g_k(n, m)| \leq \frac{2}{1 - |k|^2} |k|^{-(n+m)}.$$

Combining this bound with the identity

$$\|K(k)\|_{\mathfrak{S}_2}^2 = \sum_1^{\infty} \sum_1^{\infty} |V_n| |g_k(n, m)|^2 |V_m|$$

we obtain the claimed bound. □

Proposition 4.2 *For any $k \in \mathbb{C} \setminus \{0\}$ with $|k| < 1$,*

$$\|K(k)\|_{\mathfrak{S}_1} \leq \frac{2}{1 - |k|^2} \left(\sum_{n=1}^{\infty} |k|^{-n} |V_n|^{1/2} \right)^2,$$

Proof The matrix of $(H_0 - z)^{-1}$ is defined by

$$g_k(n, m) = \frac{k}{k^2 - 1} \left(k^{-|n-m|} - k^{-(n+m)} \right),$$

which satisfies

$$|g_k(n, m)| \leq \frac{2}{1 - |k|^2} |k|^{-(n+m)}.$$

Combining this bound with the identity

$$\|K(k)\|_{\mathfrak{S}_1} \leq \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} |V_n|^{1/2} |g_k(n, m)| |V_m|^{1/2}$$

we obtain the claimed bound. \square

5 Proof of Theorem 1.1

In this section we prove Theorem 1.1. Let us assume that V has compact support. The bound in this case implies the bound in the general case by a simple continuity argument.

As discussed in Sect. 4, the Birman–Schwinger operators $K(k)$ from (4.1) extends analytically to $\mathbb{C} \setminus \{0\}$. The same proof shows that they are not only analytic with respect to the norm of bounded operators, but even with respect to the norm in \mathfrak{S}_2 .

We will apply Corollary 2.2 to the function

$$a(k) := \det_2(1 + K(k))$$

with $\Lambda = 1/R$. Since $K(k)$ is analytic with values in \mathfrak{S}_2 , the function a is analytic. It is easy to see that assumption (2.1) is valid. Moreover, combining them with Lemma 3.1, we see that assumption (2.2) holds with

$$\ln A = \frac{1}{2} \left(\frac{2\Lambda^2}{\Lambda^2 - 1} \sum_{n=1}^{\infty} \Lambda^{2n} |V_n| \right)^2.$$

Thus, Corollary 2.2 implies that

$$\#\{j: \operatorname{Im} k_j \geq 0\} \leq \frac{1}{2 \ln \Lambda} \left(\frac{2\Lambda^2}{\Lambda^2 - 1} \sum_{n=1}^{\infty} \Lambda^{2n} |V_n| \right)^2.$$

It remains to use Lemma 3.2, which says that the $k_j + k_j^{-1}$, with $|k_j| > 1$, coincide with the eigenvalues of H , counting algebraic multiplicities. This proves Theorem 1.1.

6 Proof of Theorem 1.2

In this section we prove Theorem 1.2. Let us assume again that V has compact support.

As discussed in Sect. 4, the Birman–Schwinger operators $K(k)$ from (4.1) extend analytically to $\mathbb{C} \setminus \{0\}$. The same proof shows that they are not only analytic with respect to the norm of bounded operators, but even with respect to the norm in \mathfrak{S}_1 .

We apply Corollary 2.2 to the function

$$a(k) := \det_1(1 + K(k)) = \det(1 + K(k))$$

with $\Lambda = 1/R$. Since $K(k)$ is analytic with values in \mathfrak{S}_1 , the function a is analytic. Assumption (2.2) holds with

$$\ln A = \left(\frac{2\Lambda^2}{\Lambda^2 - 1} \sum_{n=1}^{\infty} \Lambda^n |V_n|^{1/2} \right)^2.$$

Thus, Corollary 2.2 implies that

$$\#\{j : \operatorname{Im} k_j \geq 0\} \leq \frac{1}{\ln \Lambda} \left(\frac{2\Lambda^2}{\Lambda^2 - 1} \sum_{n=1}^{\infty} \Lambda^n |V_n|^{1/2} \right)^2.$$

It remains to use Lemma 3.2, which says that the $k_j + k_j^{-1}$, with $|k_j| > 1$, coincide with the eigenvalues of H , counting algebraic multiplicities. This proves Theorem 1.2. \square

Most of the papers listed below contain results on the eigenvalues of non-selfadjoint operators. More specifically, those are the articles [1–17, 19–25, 28, 29]. The remaining references were needed for technical reasons.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

1. Abramov, A.A., Aslanyan, A., Davies, E.B.: Bounds on complex eigenvalues and resonances. *J. Phys. A* **34**, 57–72 (2001)
2. Bögli, S.: Schrödinger operator with non-zero accumulation points of complex Eigenvalues (2016). [arxiv:1605.09356](https://arxiv.org/abs/1605.09356) (Preprint)
3. Borchev, A., Golinskii, L., Kupin, S.: A Blaschke-type condition and its application to complex Jacobi matrices. *Bull. Lond. Math. Soc.* **41**, 117–123 (2009)
4. Davies, E.B.: Non-self-adjoint differential operators. *Bull. Lond. Math. Soc.* **34**(5), 513–532 (2002)
5. Davies, E.B., Nath, J.: Schrödinger operators with slowly decaying potentials. *J. Comput. Appl. Math.* **148**, 1–28 (2002)
6. Demuth, M., Katriel, G.: Eigenvalue inequalities in terms of Schatten norm bounds on differences of semigroups, and application to Schrödinger operators. *Ann. Henri Poincaré* **9**(4), 817–834 (2008)

7. Demuth, M., Hansmann, M., Katriel, G.: On the discrete spectrum of non-selfadjoint operators. *J. Funct. Anal.* **257**(9), 2742–2759 (2009)
8. Demuth, M., Hansmann, M., Katriel, G.: Eigen values of non-self adjoint operators: a comparison of two approaches. In: *Mathematical Physics, Spectral Theory and Stochastic Analysis*, Springer, pp. 107–163 (2013)
9. Enblom, A.: Estimates for eigenvalues of Schrödinger operators with complex-valued potentials (2015). [arxiv:1503.06337](https://arxiv.org/abs/1503.06337) (**Preprint**)
10. Frank, R.L.: Eigenvalue bounds for Schrödinger operators with complex potentials. *Bull. Lond. Math. Soc.* **43**(4), 745–750 (2011)
11. Frank, R.L.: Eigenvalue bounds for Schrödinger operators with complex potentials. III (2015). [arxiv:1510.03411v1](https://arxiv.org/abs/1510.03411v1) (**Preprint**)
12. Frank, R.L., Laptev, A., Lieb, E.H., Seiringer, R.: Lieb-Thirring inequalities for Schrödinger operators with complex-valued potentials. *Lett. Math. Phys.* **77**, 309–316 (2006)
13. Frank, R.L., Laptev, A., Safronov, O.: On the number of eigenvalues of Schrödinger operators with complex potentials. to appear
14. Frank, R.L., Laptev, A., Seiringer, R.: A sharp bound on eigenvalues of Schrödinger operators on the half-line with complex-valued potentials. *Spectral theory and analysis*, pp. 39–44, *Oper. Theory Adv. Appl.* **214**, Birkhäuser/Springer Basel AG, Basel (2011)
15. Frank, R.L., Sabin, J.: Restriction theorems for orthonormal functions, Strichartz inequalities and uniform Sobolev estimates (2014). [arxiv:1404.2817](https://arxiv.org/abs/1404.2817) (**Preprint**)
16. Frank, R.L., Simon, B.: Eigenvalue bounds for Schrödinger operators with complex potentials. II. *J. Spectr. Theory* (**to appear**)
17. Laptev, A., Safronov, O.: Eigenvalue estimates for Schrödinger operators with complex potentials. *Comm. Math. Phys.* **292**, 29–54 (2009)
18. Latushkin, Y., Sukhtayev, A.: The algebraic multiplicity of eigenvalues and the Evans function revisited. *Math. Model. Nat. Phenom.* **5**(4), 269–292 (2010)
19. Martirosjan, R.M.: On the spectrum of the non-selfadjoint operator $\Delta u + cu$ in three dimensional space. (Russian) *Izv. Akad. Nauk Armyan. SSR. Ser. Fiz.-Mat. Nauk* **10**(1), 85–111 (1957)
20. Martirosjan, R.M.: On the spectrum of various perturbations of the Laplace operator in spaces of three or more dimensions. (Russian) *Izv. Akad. Nauk SSSR Ser. Mat.* **24**, 897–920 (1960)
21. Murtazin, K.K.: Spectrum of the nonself-adjoint Schrödinger operator in unbounded regions. (Russian) *Mat. Zametki* **9**, 19–26. English translation: *Math. Notes* **9**(1971), 12–16 (1971)
22. Naimark, M.A.: Investigation of the spectrum and the expansion in eigenfunctions of a nonselfadjoint operator of the second order on a semi-axis. (Russian) *Trudy Moskov. Mat. Obšč.* **3**, 181–270 (1954)
23. Pavlov, B.S.: On a non-selfadjoint Schrödinger operator. (Russian) 1966 *Probl. Math. Phys.*, No. 1, *Spectral Theory and Wave Processes* (Russian) pp. 102–132 Izdat. Leningrad. Univ., Leningrad
24. Pavlov, B.S.: On a non-selfadjoint Schrödinger operator. II. (Russian) 1967 *Problems of Mathematical Physics*, No. 2, *Spectral Theory, Diffraction Problems* (Russian) pp. 133–157 Izdat. Leningrad. Univ., Leningrad
25. Safronov, O.: On a sum rule for Schrödinger operators with complex potentials. *Proc. Am. Math. Soc.* **138**(6), 2107–2112 (2010)
26. Simon, B.: *Trace Ideals and Their Applications*, 2nd edn. Amer. Math. Soc, Providence (2005)
27. Simon, B.: Notes on infinite determinants of Hilbert space operators. *Adv. Math.* **24**(3), 244–273 (1977)
28. Stepin, S.A.: Complex potentials: bound states, quantum dynamics and wave operators. *Semigroups of operators - theory and applications*, 287–297, *Springer Proc. Math. Stat.* **113**, Springer, Cham (2015)
29. Stepin, S.A.: Estimate for the number of eigenvalues of the nonselfadjoint Schrödinger operator. (Russian) *Dokl. Akad. Nauk* **455**(4), 394–397; translation in *Dokl. Math.* **89**(2), 202–205 (2014)